

WARNING

This material has been reproduced and communicated to you by or on behalf of *Charles Darwin University* in accordance with section 113P of the *Copyright Act 1968* (**Act**).

The material in this communication may be subject to copyright under the Act.
Any further reproduction or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice



Family Name					
Given Name/s					
Student Number					
Teaching Period	Semester 1, 2018				

SMA101 – Mathematics 1A	DURATION	
	Reading Time:	10 minutes
	Writing Time:	180 minutes
INSTRUCTIONS TO CANDIDATES		
<p>1 Answer all six questions.</p> <p>2 All questions are of equal value, and parts carry marks as indicated.</p> <p>3 Read ALL questions carefully.</p> <p>4 Show all working neatly in all parts. Answers without working details will attract little marks.</p> <p>5 All symbols, unless stated otherwise, have their usual meanings.</p>		
EXAM CONDITIONS		
<p><u>You may begin writing from the commencement of the examination session.</u> The reading time indicated above is provided as a guide only.</p>		
This is a CLOSED BOOK examination		
Any non-programmable calculator is permitted		
No handwritten notes are permitted		
No dictionaries are permitted		
ADDITIONAL AUTHORISED MATERIALS	EXAMINATION MATERIALS TO BE SUPPLIED	
No additional printed material is permitted	1 x 20 Page Book 1 x Scrap Paper Formula Sheet/s	

THIS EXAMINATION IS PRINTED
DOUBLE-SIDED.

THIS PAGE HAS BEEN INTENTIONALLY
LEFT BLANK.

Answer ALL questions in the Answer Booklet provided.

Marks for each question are indicated.

Question 1

- (a) Given that $f(x) = \sqrt{4 - x^2}$, find the natural domain and the range of $f(x)$. (Marks: 4)

- (b) Sketch the graph of the function:

$$y = \frac{1}{2} \ln x + 1$$

by appropriately modifying the function $y = \ln x$.

(Marks: 4)

- (c) Given that:

$$f(x) = \frac{1+x}{1-x}$$

$$g(x) = \frac{x}{1-x}$$

Find expression of $g \circ f$, and state the domain of the composition.

(Marks: 6)

- (d) Find the limit of the following:

(i) $\lim_{t \rightarrow 2} \frac{t^3 + 3t^2 - 12t + 4}{t^3 - 4t}$

(ii) $\lim_{x \rightarrow -\infty} \frac{\sqrt{3x^4 + x}}{x^2 - 8}$

(Marks: 6)

Question 2

- (a) Solve the following equalities and in each case sketch the solution on a coordinate line:

(i) $x^2 - 4x > 12$

(ii) $\frac{x}{x-3} < 4$

(Marks: 8)

- (b) A robot moves in the positive direction along a straight line so that after t minutes its distance is $s = 6t^4$ feet from the origin.

(i) Find the average velocity of the robot over the time interval $[2,4]$ minutes.

(ii) Find the instantaneous velocity of the robot at $t=2$ minutes.

(Marks: 7)

- (c) Find dy/dx of the following functions by implicit differentiation:

$$x^2y + 3xy^3 - x = 3$$

(Marks: 5)

Question 3

- (a) Using the definition of the derivative

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

of a function $f(x)$, find $f'(x)$ of $f(x) = 6x^{-3/2}$.

(Marks: 7)

- (b) Find a value of the constant k , if any, that will make the following functions continuous:

$$f(x) = \begin{cases} kx^2, & x \leq 2 \\ 2x+k, & x > 2 \end{cases}$$

(Marks: 3)

- (c) Find dy/dx of the following:

(i) $y = (5x+8)^7(1-\sqrt{x})^6$

(ii) $y = \log_{10}(\sin^2 x)$

(Marks: 5)

- (d) Evaluate the following integrals:

(i) $\int (x^2 + 3x - 1) dx$

(ii) $\int \frac{1}{1 + \sin \theta} d\theta$

(Marks: 5)

Question 4

(a) Evaluate the following definite integrals:

(i) $\int_1^4 \frac{1}{\sqrt{x}} dx$

(ii) $\int_0^\pi \sin t dt$

(Marks: 5)

(b) Express the following matrix equation as a system of linear equations:

$$\begin{bmatrix} 3 & -1 & 2 \\ 4 & 3 & 7 \\ -2 & 1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}$$

(Marks: 3)

(c) A, B, C, D and E are matrices of sizes (4×5) , (4×5) , (5×2) , (4×2) and (5×4) , respectively.

Find which of the following matrix operations are defined. For those which are defined, give the size of the resulting matrix.

- (i) BA
- (ii) $AC + D$
- (iii) $AE + B$
- (iv) $AB + B$
- (v) $E(A + B)$

(Marks: 5)

(d) Find all values of λ for which $\det(A) = 0$:

$$A = \begin{bmatrix} \lambda - 4 & 0 & 0 \\ 0 & \lambda & 2 \\ 0 & 3 & \lambda - 1 \end{bmatrix}$$

(Marks: 7)

Question 5

- (a) Find the minor M_{22} and cofactor C_{22} of the matrix:

$$A = \begin{bmatrix} 4 & -1 & 1 & 6 \\ 0 & 0 & -3 & 3 \\ 4 & 1 & 0 & 14 \\ 4 & 1 & 3 & 2 \end{bmatrix}$$

(Marks: 4)

- (b) Solve the following linear system of equations by Gaussian elimination.

$$\begin{aligned} x_1 + x_2 + 2x_3 &= 8 \\ -x_1 - 2x_2 + 3x_3 &= 1 \\ 3x_1 - 7x_2 + 4x_3 &= 10 \end{aligned}$$

(Marks: 6)

- (c) Find a unit vector that is orthogonal to both $\mathbf{u} = (1, 0, 1)$ and $\mathbf{v} = (0, 1, 1)$.

(Marks: 4)

- (d) Find a nonzero vector \mathbf{a} with terminal point $Q(3, 0, -5)$ such that \mathbf{a} is oppositely directed to vector $\mathbf{b} = (6, 7, -3)$.

(Marks: 6)

Question 6

- (a) Find the dot product $\mathbf{a} \cdot \mathbf{b}$ of the following vectors and express it in its simplest form:

(i) $\mathbf{a} = (2, 3)$ and $\mathbf{b} = (5, -7)$

(ii) $\mathbf{a} = (t, t^2)$ and $\mathbf{b} = (\cos^2 t, \frac{\sin^2 t}{t})$.

(Marks: 5)

- (b) Find $\operatorname{Re}(z)$ and $\operatorname{Im}(z)$ of:

$$z = 3e^{-\pi i}.$$

(Marks: 5)

- (c) Express the following complex expressions in the form of $a + bi$:
 $i(1 + 7i) - 3i(4 + 2i)$

(Marks: 5)

- (d) Use de Moivre's theorem to calculate the third power of:

$$1 + i$$

(Marks: 5)

Charles Darwin University
Faculty of Technology
SMA101: Formula Sheet

Calculus

● Equation of a straight line of slope m passing through a point (a, b) is : $y-b = m(x-a)$.

● If $y = 10^x$, $\log_{10} y = x$.

● $(x \pm y)^2 = x^2 \pm 2xy + y^2$

● $a^2 - b^2 = (a+b)(a-b)$

● $a^3 - b^3 = (a-b)(a^2 + ab + b^2)$

● Quadratic Eqn. $ax^2 + bx + c = 0$ has roots as $x = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a}$

● $\sin^2 x + \cos^2 x = 1$

● $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$

● $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$

● $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$

● $\frac{df}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

● $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x} = 0$

● $\frac{dx^n}{dx} = nx^{n-1}$

● $\frac{d \sin x}{dx} = \cos x$

● $\frac{d \cos x}{dx} = -\sin x$

● Product rule: $\frac{df(x)g(x)}{dx} = f(x) \frac{dg(x)}{dx} + g(x) \frac{df(x)}{dx}$

● Quotient rule: $\frac{d}{dx} \left[\frac{f}{g} \right] = \frac{g \frac{df}{dx} - f \frac{dg}{dx}}{g^2}$

● $\frac{df(g(x))}{dx} = \frac{df(g)}{dg} \frac{dg(x)}{dx}$ (Chain rule)

- $\frac{de^{\alpha x}}{dx} = \alpha e^{\alpha x}$
- Relative extremum $\frac{df(x)}{dx} = 0$
- Volume of a sphere of radius r : $V = \frac{4}{3}\pi r^3$
- Volume of a cone of radius r and height h : $V = \frac{1}{3}\pi r^2 h$.
- Area of a circle of radius r , $A = \pi r^2$
- $\frac{d \tan x}{dx} = \sec^2 x$
- $\frac{d \cot x}{dx} = -\csc^2 x$

INTEGRATION FORMULAS

DIFFERENTIATION FORMULA	INTEGRATION FORMULA	DIFFERENTIATION FORMULA	INTEGRATION FORMULA
1. $\frac{d}{dx}[x] = 1$	$\int dx = x + C$	8. $\frac{d}{dx}[-\csc x] = \csc x \cot x$	$\int \csc x \cot x dx = -\csc x + C$
2. $\frac{d}{dx}\left[\frac{x^{r+1}}{r+1}\right] = x^r \quad (r \neq -1)$	$\int x^r dx = \frac{x^{r+1}}{r+1} + C \quad (r \neq -1)$	9. $\frac{d}{dx}[e^x] = e^x$	$\int e^x dx = e^x + C$
3. $\frac{d}{dx}[\sin x] = \cos x$	$\int \cos x dx = \sin x + C$	10. $\frac{d}{dx}\left[\frac{b^x}{\ln b}\right] = b^x \quad (0 < b, b \neq 1)$	$\int b^x dx = \frac{b^x}{\ln b} + C \quad (0 < b, b \neq 1)$
4. $\frac{d}{dx}[-\cos x] = \sin x$	$\int \sin x dx = -\cos x + C$	11. $\frac{d}{dx}[\ln x] = \frac{1}{x}$	$\int \frac{1}{x} dx = \ln x + C$
5. $\frac{d}{dx}[\tan x] = \sec^2 x$	$\int \sec^2 x dx = \tan x + C$	12. $\frac{d}{dx}[\tan^{-1} x] = \frac{1}{1+x^2}$	$\int \frac{1}{1+x^2} dx = \tan^{-1} x + C$
6. $\frac{d}{dx}[-\cot x] = \csc^2 x$	$\int \csc^2 x dx = -\cot x + C$	13. $\frac{d}{dx}[\sin^{-1} x] = \frac{1}{\sqrt{1-x^2}}$	$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C$
7. $\frac{d}{dx}[\sec x] = \sec x \tan x$	$\int \sec x \tan x dx = \sec x + C$	14. $\frac{d}{dx}[\sec^{-1} x] = \frac{1}{x\sqrt{x^2-1}}$	$\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} x + C$

Linear Algebra

- If A is an invertible matrix, then:

$$A^{-1}A = AA^{-1} = I. \text{ For } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}, A^{-1} = \frac{1}{(ad-bc)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

- $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}, B = \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \\ b_{31} & b_{32} & b_{33} \end{bmatrix}$ gives

$$AB = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} + a_{13}b_{31} & a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} & a_{11}b_{13} + a_{12}b_{23} + a_{13}b_{33} \\ a_{21}b_{11} + a_{22}b_{21} + a_{23}b_{31} & a_{21}b_{12} + a_{22}b_{22} + a_{23}b_{32} & a_{21}b_{13} + a_{22}b_{23} + a_{23}b_{33} \\ a_{31}b_{11} + a_{32}b_{21} + a_{33}b_{31} & a_{31}b_{12} + a_{32}b_{22} + a_{33}b_{32} & a_{31}b_{13} + a_{32}b_{23} + a_{33}b_{33} \end{bmatrix}$$

$$\bullet \operatorname{tr}(A) = \sum_i a_{ii}$$

$$\bullet \text{ If } A \text{ is a } n \times n \text{ matrix, then: } \det(aA) = a^n \det(A) \quad \bullet \det(A^{-1}) = \frac{1}{\det(A)}$$

$$\bullet \mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta = a_1 b_1 + a_2 b_2 + a_3 b_3$$

$$\bullet \mathbf{a} \times \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \sin \theta \hat{\mathbf{n}} = \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

$$\bullet z = x + iy, \quad z^* = x - iy$$

$$\bullet z = r(\cos \theta + i \sin \theta) = re^{i\theta} : \text{polar form.}$$

$$\bullet zz^* = x^2 + y^2$$

$$\bullet (\cos \theta + i \sin \theta)^n = (\cos n\theta + i \sin n\theta)$$